

Pregnancy and Inflammation

***Emmanuel Ifeanyi Obeagu¹, Solomon Matthias Gamde² and Getrude Uzoma Obeagu³**

¹Department of Medical Laboratory Science, Kampala International University, Uganda, <https://orcid.org/0000-0002-4538-0161>

²Department of Medical Laboratory Science, Faculty of Allied Health Sciences, Bingham University Karu, Nasarawa State, Nigeria, <http://orcid.org/0000-0002-7631-8782>

³School of Nursing Science, Kampala International University, Uganda.

ABSTRACT

The concept that pregnancy is associated with immune suppression has created a myth of pregnancy as a state of immunological weakness and, therefore, of increased susceptibility to infectious diseases. A challenging question is whether the maternal immune system is a friend or a foe of pregnancy. This paper discussed the role of the immune system during pregnancy. The challenge is to better understand the immunology of pregnancy in order to deliver the appropriate treatment to patients with pregnancy complications as well as to determine public policies for the protection of pregnant women during pandemics.

Keywords: pregnancy, inflammation, cytokines.

INTRODUCTION

The idea that pregnancy is associated with immune suppression has created a myth of pregnancy as a state of immunological weakness and, therefore, of increased susceptibility to infectious diseases [1-10]. A challenging question is whether the maternal immune system is a friend or a foe of pregnancy [11-14]. In order to discuss this, question we will first review some fundamental concepts associated with the immune system and pregnancy. A fundamental feature of the immune system is to protect the host from pathogens [15]. This function depends upon the innate immune system's capacity to coordinate cell migration for surveillance and to recognize and respond to invading microorganisms. During the first trimester, NK cells, dendritic cells, and macrophages infiltrate the decidua and accumulate around the invading trophoblast cells [16]. Interestingly, depletion of immune cells, instead of helping the pregnancy, terminates the pregnancy (Hanna J et al). Thus, deletion of macrophages, NK cells, or dendritic cells (DC) has deleterious effects on placental development, implantation, or decidual formation [17]. In elegant studies, it has been shown that in the absence of NK cells, trophoblast cells are not able to reach the endometrial vascularity leading to termination of the pregnancy [18]. These studies suggest that uNK cells are critical for trophoblast invasion in the uterus. Similarly, depletion of DCs prevented blastocyst implantation and decidual formation [19]. Indeed, this study suggests that uDC are necessary for decidual formation and may affect the angiogenic response by inhibiting blood vessel maturation. More recently, Collins et al. demonstrate that uDC association, with T cell responses to the fetal "allograft," starkly contrast with their prominent role in organ transplant rejection [20]. All of these data [19], support the idea that the fetal-maternal immune interaction is more complex than the comparison to transplant allografts. Consequently, the presence of immune cells at the implantation site is not associated with a response to the "foreign" fetus but is attracted to facilitate and protect the pregnancy.

The cytokine profile during pregnancy

The definition of pregnancy as a "Th-2" or anti-inflammatory state was enthusiastically embraced and numerous studies attempted to prove and support this hypothesis [21]. This theory postulates that pregnancy is an anti-inflammatory condition [22] and a shift in the type of cytokines produced would lead to abortion or pregnancy complications. While many studies confirmed this hypothesis, a similar number of studies argued against this notion

[23]. Implantation, placentation, and the first and early second trimester of pregnancy resemble “an open wound” that requires a strong inflammatory response. The second immunological phase of pregnancy is, in many ways, the optimal time for the mother. This is a period of rapid fetal growth and development. The mother, placenta, and fetus are symbiotic, and the predominant immunological feature is induction of an anti-inflammatory state. The woman no longer suffers from nausea and fever as she did in the first stage, in part, because the immune response is no longer the predominant endocrine feature [24].

Inflammation and immune cells during implantation

As discussed above, a high level of the proinflammatory T helper (Th)-1 and cytokines (IL-6, IL-8, TNF α) characterizes early implantation [25]. These cytokines can be secreted by the endometrial cells as well as by cells of the immune system that are recruited to the site of implantation. 65–70% are uterine-specific natural killer (NK) cells [26], and 10–20% are macrophages (Mos) and 2–4 % are dendritic cells (DCs) . NK cells in human decidua have a role in regulating trophoblast invasion by the production of IL-8 and interferon-inducible protein-10 chemokines [26]. Furthermore, decidual NK cells are potent secretors of an array of angiogenic factors that induce vascular growth that is essential for the establishment of an adequate decidua [27]. DCs are a heterogeneous population of cells that initiate and coordinate the innate adaptive immune response. These cells accumulate in the pregnant uterus prior to implantation and stay in the decidua throughout pregnancy [28]. Several lines of evidence point to a pivotal role of macrophages and DCs in shaping the cytokine profile at the maternal–fetal interface [29]. Furthermore, in recent studies we showed that depletion of uterine DCs (uDCs) cells resulted in a severe impairment of implantation and led to embryo resorption [30]. However, the effect observed in our study was not related to tolerance but rather to successful decidualization. In agreement with these findings, another study showed that therapy with DCs significantly decreased the spontaneous resorption rate in a mouse model [29]. These studies, both suggest that, in addition to their involvement in the immune response, uDCs also play some trophic role in regulating the process of implantation.

CONCLUSION

Immunity in pregnancy is the result of the combination of signals and responses originating from the maternal immune system and the fetal-placental immune system. The signals originated in the placenta regulate the way the maternal immune system will behave in the presence of potential dangerous signals.

REFERENCES

1. Aluvihare VR, Kallikourdis M, Betz AG. Regulatory T cells mediate maternal tolerance to the fetus. *Nat Immunol.* 2004; 5:266–271.
2. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, Agu CC. Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria. *J Res Med Dent Sci.* 2021 Apr;9(4):145-8.
3. Okorie HM, Obeagu EI, Eze EN, Jeremiah ZA. Assessment of some haematological parameters in malaria infected pregnant women in Imo state Nigeria. *Int. J. Curr. Res. Biol. Med.* 2018;3(9):1-4.
4. Obeagu EI, Ezimah AC, Obeagu GU. Erythropoietin in the anaemias of pregnancy: a review. *Int J Curr Res Chem Pharm Sci.* 2016;3(3):10-8.
5. Obeagu EI, Hassan AO, Adepoju OJ, Obeagu GU, Okafor CJ. Evaluation of Changes in Haematological Parameters of Pregnant Women Based on Gestational Age at Olorunsogo Road Area of Ido, Ondo State, Nigeria. *Journal of Research in Medical and Dental Science.* 2021;9(12):462-4.
6. Obeagu EI, Agreen FC. Anaemia among pregnant women: A review of African pregnant teenagers. *J Pub Health Nutri.* 2023; 6 (1). 2023;138.
7. Obeagu EI, Ezimah AC, Obeagu GU. Erythropoietin in the anaemias of pregnancy: a review. *Int J Curr Res Chem Pharm Sci.* 2016;3(3):10-8.
8. Obeagu EI, Ikpenwa JN, Chukwueze CM, Obeagu GU. Evaluation of protein C, protein S and fibrinogen of pregnant women in Owerri Metropolis. *Madonna University journal of Medicine and Health Sciences ISSN: 2814-3035.* 2022 Apr 18;2(1):292-8.
9. Obeagu EI, Abdirahman BF, Bunu UO, Obeagu GU. Obstetrics characteristics that effect the newborn outcomes. *Int. J. Adv. Res. Biol. Sci.* 2023;10(3):134-43.
10. Obeagu EI, Obeagu GU, Chukwueze CM, Ikpenwa JN, Ramos GF. EVALUATION OF PROTEIN C, PROTEIN S AND FIBRINOGEN OF PREGNANT WOMEN WITH MALARIA IN OWERRI METROPOLIS. *Madonna University journal of Medicine and Health Sciences ISSN: 2814-3035.* 2022 Apr 19;2(2):1-9.
11. Bulmer JN, Pace D, Ritson A(1988). Immunoregulatory cells in human decidua: morphology, immunohistochemistry and function. *Reprod Nutr Dev.*; 28:1599–1613..

12. Okorie HM, Obeagu EI, Obarezi HC, Anyiam AF. Assessment of some inflammatory cytokines in malaria infected pregnant women in Imo State Nigeria. *International Journal of Medical Science and Dental Research*. 2019;2(1):25-36.
13. Emeka-Obi OR, Ibeh NC, Obeagu EI, Okorie HM. Evaluation of levels of some inflammatory cytokines in preeclamptic women in owerri. *Journal of Pharmaceutical Research International*. 2021;33(42A):53-65.
14. Ifeanyi O, Uzoma O, Amaeze A, Ijeko A, Felix C, Ngozi A, Nchekwubedi C, Chinenye K. Maternal expressions (serum levels) of alpha tumour necrosis factor, interleukin 10, interleukin 6 and interleukin 4 in malaria infected pregnant women based on parity in a Tertiary Hospital in Southeast, Nigeria. *Journal of Pharmaceutical Research International*. 2020 Sep 25;32 (23):35-41.
15. Mor G et al. Is the trophoblast an immune regulator?: the role of toll-like receptors during pregnancy. *Crit Rev Immunol*. 2005; 25:375-388.
16. Greenwood JD et al. Ultrastructural studies of implantation sites from mice deficient in uterine natural killer cells. *Placenta*. 2000; 21:693-702.
17. Blois SM et al. A pivotal role for galectin-1 in fetomaternal tolerance. *Nat Med*. 2007; 13:1450- 1457.
18. Hanna J et al. Decidual NK cells regulate key developmental processes at the human fetal-maternal interface. *Nat Med*. 2006; 12:1065-1074.
19. Birnberg T et al. Dendritic cells are crucial for decidual development during embryo implantation. *Am J Reprod Immunol*. 2007; 57:342
20. Collins MK, Tay CS, Erlebacher A. Dendritic cell entrapment within the pregnant uterus inhibits immune surveillance of the maternal/fetal interface in mice. *J Clin Invest*. 2009; 119:2062-2073.
21. Wegmann T (1984). Fetal protection against abortion: Is it immunosuppression or immunostimulation? *Ann Immunol*; 135D: 309-312.
22. Wegmann TG et al. Bidirectional cytokine interactions in the maternal-fetal relationship: is successful pregnancy a TH2 phenomenon? *Immunol Today*. 1993; 14:353-356.
23. Saito S, Miyazaki S, Sasaki Y. Th1/Th2 balance of the implantation site in humans. *Landes Bioscience/Springer Science*; Georgetown, Texas. 2006.
24. Romero R et al. Inflammation in preterm and term labour and delivery. *Semin Fetal Neonatal Med*. 2006; 11:317-326.
25. Jasper MJ et al. Macrophage-Derived LIF and IL1B Regulate alpha (1,2)Fucosyltransferase 2 (Fut2) Expression in Mouse Uterine Epithelial Cells During Early Pregnancy. *Biol Reprod*. 2010.
26. Manaster I, Mandelboim O. The unique properties of uterine NK cells. *Am J Reprod Immunol*. 2010; 63:434-444
27. Burke SD et al. Uterine NK cells, spiral artery modification and the regulation of blood pressure during mouse pregnancy. *Am J Reprod Immunol*. 2010; 63:472-481.
28. Scholz C et al. Glycodelin A induces a tolerogenic phenotype in monocyte-derived dendritic cells in vitro. *Am J Reprod Immunol*. 2008; 60:501-512.
29. Laskarin G, KU, Rukavina D, Thomson AW, Fernandez N, Blois SM. Antigen-presenting cells and materno-fetal tolerance: an emerging role for dendritic cells. *Am J Reprod Immunol*. 2007; 58:255-267.
30. Plaks V et al. Uterine DCs are crucial for decidua formation during embryo implantation in mice. *J Clin Invest*. 2008; 118:3954-3965.

Emmanuel Ifeanyi Obeagu, Solomon Matthias Gamade and Getrude Uzoma Obeagu (2023). Pregnancy and Inflammation. *NEWPORT INTERNATIONAL JOURNAL OF RESEARCH IN MEDICAL SCIENCES (NIJRMS)* 3 (1) :42-45.