

**NEWPORT INTERNATIONAL JOURNAL OF SCIENTIFIC AND  
EXPERIMENTAL SCIENCES  
(NIJSES)**

**Volume 4 Issue 3**

<https://doi.org/10.59298/NIJSES/2023/33.1.0151>

Page | 1

**Proximate, nutraceutical composition and antimicrobial activities of *Uvaria chamea* (Udagu) seeds and oil**

**Ogbuanu C.C.<sup>\*1</sup>, Nwagu L.N.<sup>1</sup>, Ezeh, C.N.<sup>2</sup>, Achara, N.I.<sup>3</sup>,  
Onwuatuegwu, J.T.C.<sup>4</sup>**

**<sup>1</sup>Enugu State University of Science and Technology, Department of Industrial Chemistry, Faculty of Applied Natural Sciences, P.M.B. 01660, Enugu, Nigeria**

**<sup>2</sup>University of Nigeria Enugu Campus, Centre for Environmental Management and Control**

**<sup>3</sup>Department of Biochemistry, Tansian University, Umunya**

**<sup>4</sup>Department of Microbiology, Tansian University, Umunya, Anambra State, Nigeria.**

---

**ABSTRACT**

The study investigated proximate, nutraceutical composition and Antimicrobial activities of *Uvaria chamea* (Udagu) seeds. Standard methods were used to analyze the plant seeds for the present of nutraceutical and proximate component of the plant *Uvaria chamea* (Udagu) seeds. While disk diffusion method was employed to assay the the potent of the seed oil as an antimicrobial agent. Result of the proximate revealed that *Uvaria chamea* (Udagu) seeds contains ash (10.19 %), moisture (5.2 %), crude protein (0.09 %), crude fiber (51.81 %), total fat (26.19 %) and carbohydrate (6.52 %). While nutraceutical study showed that *Uvaria chamea* (Udagu) seeds contains cardiac glycosides (0.046 %), tannins (0.2 %), saponins (0.01 %) and alkaloids (0.45 %).The proximate result implies that the plant contains essential component such as crude fiber, total fat and ash in significant quantities, and can be utilized as food and in treating several ailments. The presences of alkaloids, tannins, Cardiac glycoside, and saponins in the plant seed indicate it can be used as a medicinal source. The oil presented no antimicrobial and antibiotic activities against *Staphylococcus aureus* and *Salmonella typhi*. The results suggest that *Uvaria chamae* seed have a potential role as a new source of health-promoting diets (can be used as a dietary fiber supplement in obesity management) with high oxidative stability. And the oil is a fixed oil that could be an acceptable substitute for cooking vegetable oil that do not have antimicrobial/antibiotic activities.

Keywords: Proximate; nutraceutical; fiber; alkaloids; tannins

---

**INTRODUCTION**

Plants offer a large range of Phytochemical natural chemical compounds belonging to different molecular families which have various properties to humans as therapeutic agents that reduce the risk of cancer due to dietary fibers, polyphenol, antioxidants and anti-inflammatory effects [1]; [2]. The well-recognized father of modern medicine, Hippocrates (460-377 BC), stated "Let food be thy medicine and medicine be thy food" to predict the relationship between appropriate foods for health and their therapeutic benefits. Today phytochemicals (non-nutritive plant chemicals that have protective or disease preventative properties) in foods have been identified and new discoveries are being made

surrounding the complex benefits of phytochemicals. A combination of the words “nutrition” and “pharmaceutical,” the term nutraceutical was coined by combining the terms "Nutrition" and "Pharmaceutical" in 1989 by Dr Stephen DeFelice, Chairman of the Foundation for Innovation in Medicine, encompasses foods or food products that claim to prevent/ treat various ailments ranging from heart diseases to cancer, improve health, delay the aging process, and/or increase life expectancy [3]. Edible seed foods can be classified based on the amount of nutrients and the type of nutrients they provide for subsistence and survival and are composed of macronutrients or nutrients that are needed in large quantities (carbohydrates, proteins, fats, fiber, and water) and micronutrients or nutrients that are needed in small amount (include vitamins and minerals) in varying proportions which is essential for and nutritionally important [4]; [5], [6]; [7]. Plants capture and utilize solar energy by means of photosynthesis, thereby storing carbohydrates which serve as source of energy [8] with fiber and water, while fatty acids and *proteins, can also serve as energy sources* when they are funneled through appropriate enzymatic pathways [9]. Thus, these products serve as a source of two types of energy which can be expressed as calories. The first energy being heat which serve to maintain the body temperature. The second is free energy, which is available for work [10]. Complex macromolecule polymers of amino acids (protein) joined in peptide linkage that is very important for growth, development and maintenance that compose of 50 % of dry weight of living cells called protein that bind to other selected molecules and how their activity depends on such binding [9]; [11]. Plant fats and oils is an energy storing molecules and most important kind of lipid. They function as a structural component of membranes, aid in transportation of fat-soluble nutrients that are important for good health and as a storage form of metabolic fuel [12]; [13].

Carbohydrate nutrient are mainly derived from plants and they are polyhydroxy aldehyde or ketone or substances that yield such compounds on hydrolysis. Our body breaks down carbohydrates into glucose or blood sugar that is the main source of energy for your body's cells, tissues, and organs [14]. The primary function of the carbohydrates in living organisms is as an energy source. Many of them are easily digested by animals where they are converted back into carbon dioxide and water, with a concurrent release of energy [4]; [15]. Fiber is mainly a carbohydrate that the body can't digest but keep the digestive system healthy. A diet high in especially fiber, can help solve constipation, high cholesterol, type 2 diabetes and cardiovascular diseases, and more [16]; [17]; [18]; [19]; [20]; [21]. Fiber can help burn down trouble and stubborn fat in the body. As such fiber can help combat weight gain and keep the weight off essential for a long healthy life [22]. As a universal solvent, the moisture content (water) from food material plays an important role in living systems. It is an indispensable nutrient and involved in the metabolism of a food digestion, elimination of body wastes, transportation of nutrients, regulation of body temperature, secretion of enzymes, hormones, lubricates the joints and tissues, maintains healthy skin, and for proper digestion and other biochemical activities [23]; [24]; [25]; [26]; [27]. Edible seeds are important sources of nutrients and energy especially among the resource-poor populations where protein-energy malnutrition (PEM) has continued to hamper optimal growth and development [28]; [29] and can sustain livestock production by ensuring the availability of various sources of nutrients that are required for the formulation of animal feed [30]. Seed oils which are mainly triacylglycerols have range of physical and chemical compositional parameters that determine its applications [31]; [32]. The little or no information on the composition and utilization of the many and varied lesser seed oils indigenous to the tropics are more of problem than the real shortage of oils [33].

## MATERIALS AND METHODS

### Sample collection and preparation

The sample was ripped fruit of *Uvaria chamae* and collected from Oghe Town in Ezeagu Local Government of Enugu State on 22 July 2022. The plant was identified and authenticated by Prof. J.C. Okafor of Department of Applied Biology and Biotechnology, Enugu State University of Science and Technology. The ripped fruits were sun dried for two months and the seeds dehulled from the dried fruits and separated by the help of gravity. The whole seed was ground into powder with the aid of manual grinder and kept in a sterile plastic bag at room temperature for further studies.

### Collection of Micro-organism

The test organisms include *Staphylococcus aureus* and *Salmonella typhi* and were obtained from the Microbiology Department of Emex Medical Laboratory, Abakpa Nike Enugu. They were sub-culture on nutrient agar and kept at 37°C.

### Moisture content determination

The percent difference in weight of exactly 2g of the ground sample after 8 hours in an electric oven at 105 °C and allowed to cool down to room temperature in a desiccator containing calcium chloride for about 40 minutes [34].

Calculation

$$\% \text{ Moisture} = \frac{\text{loss in weight}}{\text{weight of sample}} \times 100$$

### Ash determination

Two grams of the ground sample was weighed into a fire-polished empty crucible and transferred into muffle furnace and heated to 550°C for 30 minutes and allowed to cool down to room temperature in a desiccator and weighed.

Calculation

$$\% \text{ Ash} = \frac{\text{Weight of ash}}{\text{Weight of sample}} \times 100$$

### Crude protein determination

The samples (1 g) was digested with 25 ml concentrated sulphuric acid, using 3 g of copper sulphate and sodium sulphate (10:1) as catalyst to convert organic nitrogen to ammonium ions. Employing macrokjeldahl method of AOAC (1999). The kjeldahl flask was inclined and heated gently at first until frothing ceased, then heated strongly with shakings, at intervals, to wash down charred particles from sides of the kjeldahl flask. Heating was continued until the mixture become clear and free from brown or black colour. The digest was allowed to cool down to room temperature and the volume made up to 100 ml with distilled water. Forty milliliters (40 ml) of 40% Sodium hydroxide was added to 20 ml of the diluted digest and the liberated ammonia is distilled with few pieces of granulated zinc and anti-bumping granules into 50 ml of 2 % boric acid receiving conical flask. The distillate was titrated with 0.01 N hydrochloric acid to a purple coloured end point (Vml).

Calculation

$$\% \text{ Nitrogen} = \frac{1.4 \times \text{Titer volume} \times \text{total volume of digest}}{1000 \times \text{weight of sample} \times \text{aliquot distilled}} \times 100$$

$$\text{Crude Protein (\%)} = \% \text{ Nitrogen} \times 6.25$$

### Crude fat determination

Five grams (5 g) of the ground sample was weighed into a thimble bag (W) and inserted into the extraction column with the condenser. The total fat content was extracted with 300 ml of the extracting solvent (n-hexane) at 60°C for six hours. After extraction, the thimble was removed and the solvent distilled out. The total fat content was placed on a pre-weighed beaker (W<sub>1</sub>) and the solvent evaporated off in a water bath and after transferred to an oven and heated for some minutes to evaporate the remaining solvent and moisture to complete dryness. It was cooled in a desiccator and weighed (W<sub>2</sub>).

The percent oil yield was calculated using

$$\text{Percent Total fat yield (\%)} = \frac{W_2 - W_1}{W} \times 100$$
$$= \frac{\text{weight of oil}}{\text{weight of Sample}} \times 100$$

### Crude fiber determination

Two grams of the defatted sample (above) obtained during fat determination was air dried and heated with 200ml of 1.25% sulphuric acid for 30 minutes and filtered. The insoluble matter was washed with boiling distilled water until the filtrate is free from acid. The insoluble matter was washed back into the flask with 200 ml of 1.25% NaOH and boiled for 30 minutes and filtered immediately. The insoluble matter was wash with boiling water until no base is detected in the filtrate. The whole insoluble matter was washed with 1% HCl and finally with boiling water until free from acid. Then it was washed twice with 95 % ethanol and three times with ether and oven-dried at 100°C to constant weight. The crucible

and its content were ash in muffle furnace at 550°C and re-weighed. The difference between the weight of ash and the weight of insoluble matter gave the weight of the crude fiber.

Calculation

$$\% \text{ Crude fiber} = \frac{\text{Weight of dried insoluble matter} - \text{weight of ash}}{\text{Weight of sample before defatting}} \times 100$$

#### Determination of total carbohydrate

The total percentage carbohydrate content was determined by difference of 100.

$\% \text{ Carbohydrate} = 100 - (\% \text{ Protein} + \% \text{ Moisture} + \% \text{ Ash} + \% \text{ Fibre})$

#### Quantitative phytochemical screening

##### Determination of total alkaloids

Twenty grams of ground *Uvariachamea* (Udagu) seeds were extracted with 500 ml of methanol: (1:1v/v) mixture and filtered with Whatmann filter paper. The filtrate was evaporated. The resultant residue was mixed with 200ml of 1 % H<sub>2</sub>SO<sub>4</sub> and partitioned with ether to remove unwanted materials. The aqueous left was then extracted five times with chloroform to obtain the alkaloid fraction. The chloroform extract was combined and concentrated to dryness. The alkaloid was weighed and the percentage was calculated with reference to the initial weight of the sample powder [35].

$$\% \text{ Alkaloids} = \frac{\text{weight of alkaloid residue} \times 100}{\text{weight of sample}}$$

##### Determination of total saponins

Twenty grams of ground sample was boiled under reflux with 300ml of 50% ethanol for 30 minutes and filtered while hot through a coarse filter paper. Activated charcoal (2 g) was added to the content to adsorb impurities (pigments), boiled and filtered while hot. The filtrate was cooled and an equal volume of acetone added to precipitate saponins. The separated saponins precipitate was dissolved in the least amount of boiling 95% alcohol and filtered while hot to remove any insoluble matter [35]. The filtrate was allowed to cool to room temperature and the precipitated saponins collected and suspended in 20ml of ethanol and filtered through a pre-weighed filter paper. The pre-weighed filter paper and the residue was transferred to a desiccator containing anhydrous calcium chloride and left to dry for 24 hrs. They were weighed with reference to the weight of sample used.

Calculation

$$\% \text{ Saponins} = \frac{\text{weight of saponins residue} \times 100}{\text{weight of sample}}$$

##### Determination of tannins

Ground (0.1g) sample was gently heated to boiling with 50 ml water for 1 hour in 100 ml conical flask and filtered. The residue was washed several times and the wash water combined with the filtrate and the solution made up to the 50 ml volume with distilled water. To 10ml of sample solution in a 50ml volumetric flask, 2.5ml of Folin - Denis reagent and 10ml of NaCO<sub>3</sub> solution were added and made to volume with distilled water and allowed to stand for 20 minutes. The optical density was measured at 760 nm using spectrophotometer [35].

Calculation

$$\text{Tannic acid (mg/ml)} = \frac{\text{absorbance of sample}}{\text{absorbance of standard} \times \text{conc. of standard}}$$

##### Determination of cardiac glycosides

One gram of the ground sample was extracted with 10ml 70% alcohol and filtered. Eight milliliters of the filtrate were diluted to a 100ml in a volumetric flask with distilled water. Eight milliliters of the diluted solution were added to 8ml of 12.5% lead acetate (to precipitate resins, tannins and pigments). The mixture was shaken well and the volume made up to 100ml with distilled water and filtered. The filtrate (50ml) was pipette into another 100ml volumetric flask and 8ml of 4.7% disodium hydrogen phosphate (Na<sub>2</sub>HPO<sub>4</sub>) solution was added to precipitate excess lead. The mixture was made up to the

volume with distilled water and stirred. This was filtered twice with filter paper. Baljets reagent (10ml) were added to 10ml of the purified filtrate and to the blank sample of 20ml of distilled water. The two solutions were incubated at room temperature for one hour (time necessary for maximum colour development). The intensity of the colour was read at 495nm using spectrophotometer. The colour was stable for several hours [35].

Calculation

$$\% \text{ glycoside} = \frac{A \times 100 \text{ g } \%}{17}$$

#### Antimicrobial assay

The antimicrobial assay of *Uvaria chamea* seed oil was performed by agar well diffusion method. Varying quantities of the oil was dissolved in Dimethyl sulfoxide (DMSO) to obtain 6.25, 12.5, 25, 50, 100 and 200 mg/ml respectively [36]. A fixed volume of Mueller Hinton agar, a standard concentration of bacteria was smeared evenly onto the surface of an agar plate. Next, disks of filter paper, each impregnated with varying concentration of the oil, were applied to the portions smeared with the organisms respectively (*Staphylococcus aureus* and *Salmonella typhi*) respectively. The plates were incubator at 37°C for 24 hours. The antimicrobial activities were observed by measuring the zone of clearance on the bacterial gene near the impregnated disc. The antibiotic activities were also performed with disc impregnated with synthetic antibiotics and zone of inhibition measured.

#### RESULTS AND DISCUSSION

Table1revealed that *Uvaria chamea* (Udagu) seeds contain some Nutraceutical composition which also plays their own vital role. The result showed that the pant seed contains0.046 % cardiac glycoside, 0.2 % tannins, 0.01 % saponin and 0.45 % alkaloids. This property is also due to the plant genetic makeup.

**Table 1: Nutraceutical composition of *Uvaria chamea* (Udagu) seeds**

| S/N | Parameter         | Composition (%) |
|-----|-------------------|-----------------|
| 1   | Cardiac glycoside | 0.046           |
| 2   | Tannins           | 0.2             |
| 3   | Saponins          | 0.01            |
| 4   | Alkaloids         | 0.45            |

The findings of the study showed that the seeds of *Uvaria chamea* (Udagu) plant contain phytochemicals such as alkaloids, tannins, Cardiac glycoside, and saponins. Evidence from the study indicated that the concentration of the parameters varies. The presences of alkaloids, tannins, Cardiac glycoside, and saponins in the plant seed indicate that the plant can be used as medicinal source. The findings also corroborate with the findings of [37] who in their study said that tannin content of the thirty browse plants was 0.13 %-1.07 %and the small quantity in *tannin* content of the *Uvaria chamea* seeds may likely suggests that the seed is not rich in tannins. The percent ash, moisture, crude protein, crude fiber, total fat and carbohydrate contents of the *Uvaria chamea* (Udagu) seeds were analyzed using simple chemical methods respectively. Result in Table2 shows that the *Uvaria chamea* (Udagu) seeds contain 10.19 % ash, 5.2 % moisture, 51.81 % crude fiber, 0.09 % crude protein, 26.19 % total fat and 6.48 % carbohydrate. This result implies that the plant contains essential component such as crude fiber (51.81 %), total fat (26.19 %) and ash (10.19 % ash), and can be utilized as food and in treating several ailments. This result may also be attributed to genetic composition of the plant The *Uvaria chamea*(Annonaceae)which has been reported to have potent activity against lymphocytic leukemia in mice [38], anti-malarial activity [39], it also has the anticarcinogenic [40]; [41], exhibited significant in vitro cytotoxicity against the KB cancer cell line [42], has antidiabetic activity [43-50] has antihemolytic properties [50-57], has antimicrobial activity [58-65] has high content of phenolic compounds and high antioxidant activity [46].

**Table 2: Proximate Determination and percent content of *Uvaria chamea* (Udagu) seeds**

| S/N | (%) Parameter | Composition (%) |
|-----|---------------|-----------------|
| 1   | Ash           | 10.19           |
| 2   | Moisture      | 5.2             |
| 3   | Crude protein | 0.09            |
| 4   | Crude fiber   | 51.81           |
| 5   | Total fat     | 26.19           |
| 6   | Carbohydrate  | 6.52            |

The proximate content of the plant with their varying concentration such as that of carbohydrate (6.52 %) and crude fiber (51.81 %) indicates that the plant can be a good source of energy and fiber for animals. The findings of the study that the seed is not a good source of crude protein. Evidence from the study also showed that *Uvaria chamea* (Udagu) seeds also is a good source of oil. The most interesting components. The antimicrobial activity of the *Uvaria chamea* seed oil was performed using disc diffusion method and Mueller Hinton agar. The results are presented in Table 3.

**Table 3: Antibacterial activity of *Uvaria chama* seed oil against pathogenic microbes.**

| Microbes                     | <i>Uvarian chamea</i> seed oil |
|------------------------------|--------------------------------|
| <i>Staphylococcus aureus</i> | No activity                    |
| <i>Salmonella typhi</i>      | No activity                    |

There were no antimicrobial activities against the two microbes (*Staphylococcus aureus* and *Salmonella typhi*) used in this study.

**Table 4: Antibacterial-zone inhibition (mm) activity of antibiotic disc on the test organism.**

| Test organism     | Zone of inhibition (mm) |     |     |     |     |    |     |     |     |     |  |
|-------------------|-------------------------|-----|-----|-----|-----|----|-----|-----|-----|-----|--|
| Gram +ve bacteria | APX                     | Z   | AM  | R   | CPX | S  | SXT | E   | PEF | CN  |  |
| <i>S. aureus</i>  | 0.0                     | 19  | 0.0 | 0.0 | 22  | 18 | 0.0 | 15  | 18  | 0.0 |  |
| Gram -ve bacteria | CH                      | OFX | SP  | SXT | PEF | S  | CPX | AM  | AU  |     |  |
| <i>E. coli</i>    | 20                      | 17  | 17  | 20  | 16  | 21 | 18  | 0.0 | 0.0 |     |  |

**Key:**

Gram +ve = APX → Ampiclox; Z → Zinnacef; AM → Ampicilin; R → Recephin; CPX → Cyproflacin; S → Streptomacin; SXT → Septrin; E → Erytromycin; PEF → Reflacin; CN → Gentamycin.

Gram -ve = CH → Chlorophenicol; OFX → Tarvid; SXT → Septrin; PEF → Reflacin; S → Streptomacin; CPX → Cyproflacin; AM → Ampicilin; AU → Augmentin

The results obtained shows that Zinnacef, Cyproflacin, Streptomacin, Erytromycin and Reflacin were potent against the gram positive test organisms. While Chlorophenicol, Tarvid, Septrin, Reflacin, Streptomacin, Cyproflacin and Augmentin have *potent* against gram negative test bacteria.

**CONCLUSION**

Evidence from the study clearly indicated that *Uvaria chamea* (Udagu) seeds have some potential medicinal properties (alkaloid, tannins, saponin and cardiac glycoside). This phytochemical and other founds in the plant makes the *Uvaria chamea* (Udagu) seeds a potential medicinal plant. The fiber and oil content are significant and hence, it can be accepted that the plant seed can be harnessed for nutrition, obesity management and industrial purposes. The results of the present investigation clearly demonstrate that the oil of *Uvaria chamea* seeds have no activity against the two test organisms (*S. aureus* and *E. coli*) respectively. This work justifies the standard antibiotics have potent against the two

test organisms (*S. aureus* and *S. typhi*) respectively with exception of ampiclox, ampicillin, septrin and gentamycin that have no potent against gram positive bacteria's, while for gram negative bacteria's only ampicillin and augmentin that have no potent.

#### Suggestion for further study

Study should be carried out on the effect of *Uvaria chamea* (Udagu) seeds oil for human consumption.

#### REFERENCES

1. Hervé, Z., Charles, K., Anoubilé, B., Janat, M.B. and Yves, A.B. (2008). Phytochemical Screening and Determination of Flavonoids in *Secamoneafzelii* (Asclepiadaceae) Extracts. *Afr. J. Pure Appl. Chem.*, 2(8): 80-82.
2. Kinderley, D. (2006). *Nutrition for life*. Lark and Deen Publishers, UK, p 213.
3. Muredzi, P. (2013). *Food is Medicine – An Introduction to Nutraceuticals.*; LAP Lambert Academic Publishing; ISBN 978-3659437106.
4. Morris, A.L and Mohiuddin, S.S. (2022). Biochemistry, Nutrients. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing.
5. Samtiya, M., Aluko, R.E. and Dhewa, T. (2020). Plant food anti-nutritional factors and their reduction strategies: an overview. *Food Prod Process and Nutr* 2, 6.
6. Abiola Oso, A. and Omotayo Ashafa, A.. (2021).. Nutritional Composition of Grain and Seed Proteins. Grain and Seed Proteins Functionality [Working Title]. doi: 10.5772/intechopen.97878
7. Chen, C., Chaudhary, A. and Mathys, A. (2021). Nutrient Adequacy of Global Food Production. *Front. Nutr.* 8:739755. doi: 10.3389/fnut.2021.739755.
8. Janssen, P.J.D., Lambreva, M.D., Plumeré, N., Bartolucci, C., Antonacci, A., Buonasera, K., Frese, R.N., Scognamiglio, V. and Rea, G. (2014). Photosynthesis at the forefront of a sustainable life. *Front. Chem.* 2:36. doi: 10.3389/fchem.2014.00036
9. Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K. and Walter, P. (2002). *Molecular Biology of the Cell*. 4th edition. New York: Garland Science. How Cells Obtain Energy from Food. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK26882/>
10. Popson, M.S., Dimri, M and Borger, J. (2022). Biochemistry, Heat and Calories. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; Available from: <https://www.ncbi.nlm.nih.gov/books/NBK538294/>
11. Nelson, D. L. and Cox, M. M. (2012). *Lehninger Principle of Biochemistry*, fourth edition:
12. NIH, National Institute of General Medical Sciences (NIGMS). (2013). The biology of fats in the body. *Science Daily*. Retrieved March 2, 2023 from [www.sciencedaily.com/releases/2013/04/130423102127.htm](http://www.sciencedaily.com/releases/2013/04/130423102127.htm)
13. El-Zayat, S.R., Sibaii, H. and El-Shamy, K.A. (2019). Physiological process of fat loss. *Bull Natl Res Cent* 43, 208. <https://doi.org/10.1186/s42269-019-0238-z>
14. Saha, S.K. and Pathak, N.N. (2021). Carbohydrates. In: *Fundamentals of Animal Nutrition*. Springer, Singapore. [https://doi.org/10.1007/978-981-15-9125-9\\_6](https://doi.org/10.1007/978-981-15-9125-9_6)
15. Holesh, J.E., Aslam, S. and Martin, A. (2022). Physiology, Carbohydrates. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing.
16. Ma, W., Nguyen, L.H., Song, M., Jovani, M., Liu, P.H., Cao, Y., Tam, I., Wu, K., Giovannucci, E.L., Strate, L.L. and Chan, A.T. (2019). Intake of dietary fiber, fruits and vegetables, and risk of diverticulitis. *American Journal of gastroenterology*. 114(9): 1531
17. Jesch, E.D. and Carr, T.P. (2017). Food ingredients that inhibit cholesterol absorption. *Preventive nutrition and food science*; 22(2): 67.
18. Brown, L, Rosner, B., Willett, W.W. and Sacks, F.M. (1999). Cholesterol-lowering effects of dietary fiber: a meta-analysis. *The American journal of clinical nutrition*. ;69(1): 30-42.
19. McKeown, N.M., Meigs, J.B., Liu, S., Wilson, P.W., Jacques, P.F. (2002). Whole-grain intake is favorably associated with metabolic risk factors for type 2 diabetes and cardiovascular disease in the Framingham Offspring Study. *The American journal of clinical nutrition*.;76(2): 390-8.
20. Hu, Y., Ding, M., Sampson, L., Willett, W.C., Manson, J.E., Wang, M., ROSNER, B., Hu, F.B. and Sun, Q. (2020). Intake of whole grain foods and risk of type 2 diabetes: results from three prospective cohort studies. *BMJ* 370
21. Lattimer, J.M. and Haub, M.D. (2010). Effects of Dietary Fiber and Its Components on Metabolic Health. *Nutrients*, 2: 1266-1289.

22. Hairston, K. G., Vitolins, M. Z., Norris, J. M., Anderson, A. M., Hanley, A. J. and Wagenknecht, L. E. (2012). Lifestyle Factors and 5-Year Abdominal Fat Accumulation in a Minority Cohort: The IRAS Family Study. *Obesity*, 20(2), 421–427.
23. Olaseni, A., Moses, M., Mariam Oyefunke, O. & Saheed, S. (2021). 11 Dietary nutrients, antinutritional factors, and valorization of food waste. In O. Ijabadeniyi (Ed.), *Food Science and Technology: Trends and Future Prospects* (pp. 281-322). Berlin, Boston: De Gruyter.
24. Millard-Stafford, M., Wendland, D.M., O'Dea, N.K. and Norman, T.L. (2012). Thirst and hydration status in everyday life. *Nutr Rev.*;70 Suppl 2:S147-51.
25. Hopkins, W.G. and Huner, N.P.A. (2009). *Introduction to Plant Physiology*, 4th Edition. John Wiley and Sons, Inc., Hoboken, USA.
26. Salisbury, F.B. and Ross, C.W. (1992). *Plant Physiology*. Wadsworth Publishing Company, Belmont - California.
27. Taiz, L. and Zeiger, E. (2002 and 2010). *Plant Physiology*, 3rd and 5th Edition. The Benjamin Cummings Publishing Company, Redwood City - California.
28. Perumal, S., Klaus, B., Harinder, P.S.M. (2001). Chemical composition, protein fractionation, essential amino acid potential and antimetabolic constituents of an unconventional legume, Gila bean (*Entadaphaseoloides Merrill*) seed kernel. *Journal of Science of Food and Agriculture* 82: 192-202.
29. Escudero, N.L., Zirulnik, F., Gomez, N.N., Mucciarelli, S.I., Gimenez, M.S. (2006). Influence of a protein concentrate from *Amaranthus cruentus* seeds on lipid metabolism. *Experimental Biology and Medicine* 231: 50-59.
30. Amole, T. , Augustine, A. , Balehegn, M. , & Adesogoan, A. T. (2022). Livestock feed resources in the West African Sahel. *Agronomy Journal*, 114, 26–45.
31. Frank D.G., *The chemistry of oils and fats: Sources, composition, properties and uses*. Black well publishing Ltd, pp 1-2, 2004.
32. Beare-Rogers J.L., Trans and positional isomers of common fatty acids. In HH Draper (ed.) *Advances in Nutritional Research*. Vol. 5 Plenum Press, New York, pp 171-200, 1983.
33. Anon A., *World fats and oil report*. *Journal of American Oil Chemists' Society*, 1987; 64: 1058 – 1085.
34. Ogbuanu C. C., Chime C. C. and Nwagu L. N. (2015). Physiochemical and fatty acid analysis of *Virescens*(Ojukwu) oil and *Nigrescens*(ordinary) palm oil of *Eleaisguineensis*. *AJFS* 9(7): 400-405
35. Evans, W. C and Trease.L (1999). *Trease and Evans Pharmacognosy 14th Edition W. B. Saunders Company Limited*, New York pp1-340.
36. Maragathavalli, S., Brindha, S., Kaviyarasi, N.S., Annadueai, B. and Gargwar, S.K. (2012). Antimicrobial activity of leaf extract of Neem (*Azadirachta indica Linn*). *International Journal of Science and Nature*, 3(1): 110-113.
37. Kökten, K. , Koçak, A. , Bağcı, E. , Akçura, M. and ÇelİK, S. (2010). Tannin, protein contents and fatty acid compositions of the seeds of several *Vicia L.* species from Turkey. *Grasas y Aceites (Sevilla)* 61(4): 404-408.
38. Jolad, S.D., Hoffmann, J.J., Schram, K.H. and Cole, J.R. (1982). Uvaricin, a new antitumor agent from *Uvariaacuminata*(Annonaceae). *J. Org. Chem.* 47: 3151-3153.
39. Nkunya, M., Moshi, M., Joseph, C. and Innocent, E. (1991). In vitro antibacterial activities of extracts and compounds from *Uvariascheffleri*. *Pharmaceutical Biology*, 1744-5116. 42(4): 267-273.
40. Popoola, T.D., Guetchueng, S.T., Ritchie, K.J. *et al.* (2021). Potent Nrf2-inducing, antioxidant, and anti-inflammatory effects and identification of constituents validate the anti-cancer use of *Uvariachamae* and *Olaxsubscorpioidea*. *BMC Complement Med Ther* 21, 234
41. Popoola TD, Awodele O, Babawale F, Oguns O, Onabanjo O, Ibang I, Godwin H, Oyeniyi T, Fatokun AA, Akinloye O. (2019). Antioxidative, antimitotic, and DNA-damaging activities of *Garcinia kola* stem bark, *Uvariachamae* root, and *Olaxsubscorpioidea* root used in the ethnotherapy of cancers. *J Basic Clin Physiol Pharmacol*;31(3):/j/jbcpp.2020.31.issue-3/jbcpp-2019-0073/jbcpp-2019-0073.xml.
42. Véronique Eparvier, Van Hung Nguyen, Odile Thoison, Françoise Guéritte, Marie-Thérèse Martin and Thierry Sévenet (2006). Cytotoxic Monotetrahydrofuran Acetogenins from *Disepalumplagioneurum*. *Journal of Natural Products* 69(9):1289-94.
43. Emordi, J.E., Agbaje, J.E., Oreagba, I. A and Iribhogbe, O. I (2018). Antidiabetic Effects of the Ethanolic Root Extract of *U. chamae* P. Beauv (Annonaceae) in Alloxan-Induced Diabetic Rats:



- A Potential Alternative Treatment for Diabetes Mellitus. *Advances in Pharmacological Sciences*, 2018: 1-13
44. Avaligbe, C.T., Joachim, D., Gbenou, D. S., Salomé Kpoviessi, Georges C. Accrombessi, M., Moudachirou and M. Gbeassor (2012). Antihemolytic Properties of Extracts of Six Plants Used in the Traditional Treatment of Sickle Cell Disease in Benin. *Journal of Applied Pharmaceutical Science* 02 (03): 08-13.
  45. Ebi, G, C., Ifeanacho C.J. and Kamalu T.N. (1999). Antimicrobial properties of *U. chamae* stem bark. *Fitoterapia*, 70(6): 621-624.
  46. Kone, M., Toure, A., Ouattara, K., Coulibaly, A. (2015). Phytochemical Composition, Antioxidant and Antibacterial Activities of Root of *U. chamae* P. Beauv. (Annonaceae) Used in Treatment of Dysentery in North of Côte d'Ivoire. *International Journal of Pharmacognosy and Phytochemical Research*; 7(6); 1047-1053.
  47. Okechukwu, P. U., Okwesili, F. N., Parker, E. J., Abubakar, B., Emmanuel, C. O., & Christian, E. O. (2013). Phytochemical and acute toxicity studies of *Moringa oleifera* ethanol leaf extract. *International Journal of Life Science BiotechNology and Pharma Research*, 2(2), 66-71.
  48. Odo, C. E., Nwodo, O. F., Joshua, P. E., Ugwu, O. P., & Okonkwo, C. C. (2013). Acute toxicity investigation and anti-diarrhoeal effect of the chloroform-methanol extract of the seeds of *Persea americana* in albino rats. *journal of pharmacy research*, 6(3), 331-335.
  49. Adonu Cyril, C., Ugwu, O. P. C., Esimone Co, O., Bawa, A., Nwaka, A. C., & Okorie, C. U. (2013). Phytochemical analyses of the menthanol, hot water and n-hexane extracts of the aerial parts of *cassytha filiformis* (Linn) and leaves of *cleistopholis patens*. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 4, 1143-1149.
  50. Orji, O. U., Ibiam, U. A., Aja, P. M., Ugwu, P., Uraku, A. J., Aloke, C., ... & Nwali, B. U. (2016). Evaluation of the phytochemical and nutritional profiles of *Cnidoscopus aconitifolius* leaf collected in Abakaliki South East Nigeria. *World Journal of Medical Sciences*, 13(3), 213-217.
  51. Offor, C. E., Ugwu, P. C., Okechukwu, P. M., & Igwenyi, I. O. (2015). Proximate and phytochemical analyses of *Terminalia catappa* leaves. *European Journal of Applied Sciences*, 7(1), 09-11.
  52. Nwali, B. U., Egesimba, G. I., Ugwu, P. C. O., & Ogbanshi, M. E. (2015). Assessment of the nutritional value of wild and farmed *Clarias gariepinus*. *International Journal of Current Microbiology and Applied Sciences*, 4(1), 179-182.
  53. Afiukwa, C. A., Igwenyi, I. O., Ogah, O., Offor, C. E., & Ugwu, O. O. (2011). Variations in seed phytic and oxalic acid contents among Nigerian cowpea accessions and their relationship with grain yield. *Continental Journal of Food Science and Technology*, 5(2), 40-48.
  54. Aja, P. M., Okechukwu, P. C. U., Kennedy, K., Ibere, J. B., & Ekpono, E. U. (2017). Phytochemical analysis of *Senna occidentalis* leaves. *IDOSR J Appl Sci*, 2(1), 75-91.
  55. Igwenyi, I. O., Isiguzo, O. E., Aja, P. M., Ugwu Okechukwu, P. C., Ezeani, N. N., & Uraku, A. J. (2015). Proximate composition, mineral content and phytochemical analysis of the African oil bean (*Pentaclethra macrophylla*) seed. *American-Eurasian J Agric Environ Sci*, 15, 1873-1875.
  56. Orji, O. U., Ibiam, U. A., Aja, P. M., Ugwu, P., Uraku, A. J., Aloke, C., ... & Nwali, B. U. (2016). Evaluation of the phytochemical and nutritional profiles of *Cnidoscopus aconitifolius* leaf collected in Abakaliki South East Nigeria. *World Journal of Medical Sciences*, 13(3), 213-217.
  57. Offor, C. E., Ugwu, P. C., Okechukwu, P. M., & Igwenyi, I. O. (2015). Proximate and phytochemical analyses of *Terminalia catappa* leaves. *European Journal of Applied Sciences*, 7(1), 09-11.
  58. Afiukwa, C. A., Ugwu, O. P., Ebenyi, L. N., Oketa, H. A., Idenyi, J. N., & Ossai, E. C. (2013). Phytochemical analysis of two wild edible mushrooms, *Auricularia polytricha* and *Pleurotus ostreatus*, common in Ohaukwu area of Ebonyi state, Nigeria. *Res J Pharm Biol Chem Sci*, 4(2), 1065-70.
  59. Chukwuemeka, I. M., Udeozo, I. P., Mathew, C., Oraekwute, E. E., Onyeze, R. C., & Ugwu, O. P. C. (2013). Phytochemical analysis of crude ethanolic leaf extract of *Morinda lucida*. *Int. J. Res. Rev. Pharm. Appl. Sci*, 3(4), 470-475.
  60. Udeozo, I. P., Nwaka, A. C., Ugwu, O. P., & Akogwu, M. (2014). Anti-inflammatory, phytochemical and acute toxicity study of the flower extract of *Newbouldia laevis*. *Int J Curr Microbiol App Sci*, 3(3), 1029-35.

61. Afiukwa, C. A., Ugwu Okechukwu, P. C., Ebonyi, L. N., Ossai, E. C., & Nwaka, A. C. (2013). Phytochemical analysis of three wild edible mushrooms, coral mushroom, *Agaricus bisporus* and *Lentinus sajor-caju*, common in Ohaukwu Area of Ebonyi State, Nigeria. *International Journal of Pharmaceutics*, 3(2), 410-414.
62. PC, U. O., & Amasiorah, V. I. (2020). The effects of the crude ethanol root extract and fractions of *Sphenocentrum jollyanum* on hematological indices and glycosylated haemoglobin of streptozotocin-induced diabetic albino rats. *INOSR Scientific Research*, 6(1), 61-74.
63. Ikechukwu, A. A., Ibiam, U. A., Okechukwu, P. U., Inya-Agha, O. R., Obasi, U. O., & Chukwu, D. O. (2015). Phytochemistry and acute toxicity study of *Bridelia ferruginea* extracts. *World J. Med. Sci*, 12(4), 397-402.
64. Igwenyi, I. O., Dickson, O., Igwenyi, I. P., Okechukwu, P. C., Edwin, N., & Alum, E. U. (2015). Properties of Vegetable Oils from Three Underutilized Indigenous Seeds. *Global Journal of Pharmacology*, 9(4), 362-365.
65. Ibiam, U. A., Alum, E. U., Aja, P. M., Orji, O. U., Nwamaka, E. N., & Ugwu, O. P. C. (2018). Comparative Analysis of Chemical Composition of *Buchholzia Coriacea* Ethanol Leaf-Extract, Aqueous and Ethylacetate Fractions. *INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES*, 5(7), 6358-6369.

**CITE AS: Ogbuanu C.C., Nwagu L.N., Ezech, C.N., Achara, N.I., Onwuatuegwu, J.T.C. (2023). Proximate, nutraceutical composition and antimicrobial activities of *Uvaria chamea* (Udagu) seeds and oil. NEWPORT INTERNATIONAL JOURNAL OF SCIENTIFIC AND EXPERIMENTAL SCIENCES, 4(3): 1-10. <https://doi.org/10.59298/NIJSES/2023/33.1.0151>**